

Research Article**The first continuous plankton sampling by VOR (Towed Oceanographic Vehicle) in southeastern and southern Brazil waters**Márcio Silva Tamanaha¹, Dyegho Moraes Costa-Gama Cunha¹ & Charrid Resgalla Junior¹¹Centro de Ciências Tecnológica da Terra e do Mar (CTTMar)

Universidade do Vale do Itajaí, Itajaí, Santa Catarina, Brazil

Corresponding author: Márcio Silva Tamanaha (marcio.tamanaha@univali.br)

ABSTRACT: Plankton sampling with the Continuous Plankton Sampling survey has been conducted since 1938 in the Northern Atlantic Ocean and North Sea, where in both cases historical records have documented climate change. Plankton sampling with the VOR (Towed Oceanographic Vehicle) on industrial fishing boats was performed in May-June of 2013 off the southeastern and southern coasts of Brazil (between 23°00'S-44°21'W and 31°15'S-50°06'W). The speed of VOR towing onboard the fishing vessels was 8-10 knots to 10-12 m of depth. The size of mesh silk used was 250-270 µm. The total of taxa is 73 phytoplankton, 26 zooplankton, 10 protozooplankton (Ciliophora) and ichthyoplankton (fish eggs) were recorded. The dominant phytoplankton groups detected were diatoms and dinoflagellates, although cyanobacteria were also abundant in the southeast region. The zooplankton with a size class of less than 2 mm was the most frequent. A positive correlation was observed between herbivorous zooplankton and phytoplankton in the coastal waters.

Keywords: continental shelf, ship of opportunity, VOR, plankton, Brazil.

Primer muestreo continuo de plancton con VOR (Vehículo Oceanográfico Remolcado) en aguas del sureste y sur de Brasil

RESUMEN: El muestreo de plancton con el Continuous Plankton Recorder (CPR) se ha efectuado desde el año 1938 en el Océano Atlántico Norte y Mar del Norte, donde en ambos casos el registro histórico ha documentado el cambio climático. El muestreo de plancton con el VOR (Vehículo Oceanográfico de Remolque), empleado por la flota pesquera industrial se efectuó en mayo-junio de 2013 en la costa sur y sureste de Brasil (23°00'S-44°21'W y 31°15'S-50°06'W). La velocidad de arrastre del VOR, utilizada por las embarcaciones pesqueras fue de 8-10 nudos a 10-12 m de profundidad, utilizando un tamaño de red de 250-270 µm. Se registró un total de 73 taxa de fitoplancton, 26 de zooplancton, 10 de protozoarios e ictioplancton (huevos de peces). Los grupos dominantes de fitoplancton fueron diatomeas y dinoflagelados, aunque las cianobacterias fueron también abundantes en la región sudeste. El zooplancton con un tamaño de clase <2 mm fue el más frecuente. Se observó una correlación positiva entre el zooplancton herbívoro y fitoplancton en aguas costeras.

Palabras clave: plataforma continental, barco de oportunidad, VOR, plancton, Brasil.

INTRODUCTION

Study of marine plankton using the CPR (Continuous Plankton Recorder) began in the North Sea in 1938 (Lucas, 1941), with the methods being later standardized, in 1960 (Colebrook, 1960). Since then, the methodology has not been significantly changed. The majority of studies have focused on the North Atlan-

tic, North Sea and North Pacific (Warner & Hays, 1994; John *et al.*, 2002; Lindley & Batten, 2002; Batten *et al.*, 2003a; Richardson *et al.*, 2006; Head & Pepin, 2010a, 2010b). In the southern hemisphere, sampling began in 2001 (Hosie *et al.*, 2003; Hunt & Hosie, 2003) with emphasis on describing the variation of plankton between the Australian (60° to 160°E and 48°S) coastal and oceanic zone. In the last decades, studies have been

conducted using historical data of CPR in relation to climate change (Edwards *et al.*, 2001; Head & Pepin, 2010a), and phenology of the plankton (Edwards & Richardson, 2004).

Different lines of research associated with the planktonic community can be conducted through the use of Continuous Plankton Sampling (CPS), which facilitates obtaining data in regions which until then, had not been studied on the same spatial-temporal scale. For the Brazilian coast, the studies of plankton were always based on sampling with traditional sampling methods, using plankton nets and bottles (Alvarino, 1980; Moser & Giancesella-Galvão, 1997), involving different regions of the coast in the north (Araujo & Ribeiro, 2005), northeast (Klein, 1977; Gomes *et al.*, 1989), southeastern (Muxagata & Montú, 1999), and southern (Resgalla Jr. *et al.*, 2001; Resgalla Jr., 2011), and in all cases, the samplings were conducted by scientific cruises, with limited duration and geographical coverage.

The opportunity to obtain oceanographic data continuously, using the VOR (Towed Oceanographic Vehicle - from the Portuguese Veículo Oceanográfico de Reboque) developed by Faccin *et al.* (2014), used in different fishing boats of the industrial fleet (opportunity vessels), offers new perspectives for the knowledge of planktonic groups on the Brazilian continental shelf. Associated with this, the use of fishing boats means that the oceanographic parameters can be easily collected in marine areas of greater economic interest and ecological importance. This study therefore presents the initial results obtained by continuous plankton sampling with the use of the VOR, towed by vessels of the industrial fishing fleet in the southeastern and southern regions of Brazil. The aim of this study is to assess plankton community distribution through continuous plankton monitoring by VOR in different Brazilian Coastal Zones and their relationship with local hydrological characteristics.

MATERIALS AND METHODS

Study area

The Continental Shelf of the South Atlantic (Southwestern Atlantic Shelf - SWAS) comprises the southern and southeastern regions of Brazil involving the Southern Brazilian Bight (SBB - $\sim 22^{\circ}00' - 28^{\circ}30'S$) and Southern Subtropical Shelf (SSS - $\sim 28^{\circ}30' - 33^{\circ}75'S$) (Piola *et al.*, 2000; Lopes *et al.*, 2006; Palma *et al.*, 2008). The prevalent water masses in the SBB are Tropical Water (TW - $T > 18.5^{\circ}C$ and $S > 36.4$) of the Brazil Current on the surface, South Atlantic Central Water (SACW - $T < 20^{\circ}C$ and $S < 36.4$) in the sub-

surface (200-500 m), and Coastal Water (CW) (50 m) influenced by different river inflows to the Brazilian coastal zone. In the SSS, the Falkland Current is also present, especially in the colder months, together with the Subantarctic Water mixed with the Rio de La Plata, forming the La Plata Front (Piola *et al.*, 2000; Palma *et al.*, 2008).

Sampling and data acquisition

The VOR (Faccin *et al.*, 2014) operates with a silk mesh (250 and 270 μm), at trawling depths between 10 and 12 m and speeds between 8 and 10 knots (15-18.5 $km h^{-1}$). Sampling was conducted from May-June 2013 on industrial fishing fleet in the region of Itajaí (southern Brazil). Two trawlers were used: the Mtanos Seif operates with purse-seine fishing targeting *Sardinella brasiliensis* and the Nicassio da Costa operates with simple trawling targeting *Umbrina canosai* and *Prionotus punctatus*. The trawls (transects) were performed on the continental shelf of the southeastern and southern regions of Brazil, between the states of Rio de Janeiro and Rio Grande do Sul (Table 1, Fig. 1). In all the trips, the VOR was operated by scientific observers on board, with the aid of the crew.


After each trawl, the mesh of the plankton collection unit (K7) was rewound and wrapped in plastic bags containing 4% formaldehyde buffered with Potassium and Sodium tetraborate. The trawl courses were determined using GPS Archer software Navicomputer[®], and the data on sea surface temperature between the areas sampled were obtained from satellite data from the NASA website for the same periods (http://gdata1.sci.gsfc.nasa.gov/daac-bin/G3/gui.cgi?instance_id=ocean_month).

Plankton analysis

In the laboratory, the meshes were sectioned every 5 nautical miles. Different analysis methods were used for the phytoplankton, protozooplankton and zooplankton, according to Batten *et al.*, (2003b) and Richardson *et al.* (2006). Qualitative and quantitative analyses of the phytoplankton were performed under a CPR microscope (platinum glass - 150x200 mm). At least 30 counting fields were used, with cross-sectional identification with 625x magnification and a field area to mesh area ratio 1/8000. The protozooplankton and zooplankton <2 mm were analysed by scanning with 65x magnification, and a field area to mesh area ratio of 1/50. After the work with the phytoplankton and the protozooplankton, zooplankton >2 mm were removed from filter mesh, counted and identified in a Bogorov chamber in their totality. All groups were identified to the lowest possible taxon; for the zooplankton, there was a prevalence of copepods.

Table 1. Details of VOR transects conducted during fishing vessels cruises in May and June 2013.

Transect	Vessel (fishing boats)	Sampling period	Latitud (S)	Longitud (E)	Distance (nautical miles)
SSB1	<i>Mtanos Seif</i>	May 02, 2013	23.00-24.56	44.21-44.55	30.00
SSB2	<i>Mtanos Seif</i>	May 03, 2013	24.56-25.31	47.03-47.31	40.00
SSB3	<i>Mtanos Seif</i>	May 03, 2013	26.24-26.25	48.26-47.58	20.00
SSB4	<i>Nicássio Costa</i>	June 18, 2013	27.03-27.17	48.22-47.57	25.00
SSS1	<i>Nicássio Costa</i>	June 28, 2013	30.45-30.26	49.47-49.13	20.00
SSS2	<i>Nicássio Costa</i>	June 28, 2013	31.15-31.32	49.59-50.06	35.00

**Figure 1.** VOR map towing from industrial fishing boats. Highlight of the transects of the southeast and south regions of Brazil (solid line: SSB1, SSB2, SSB3, SSB4, SSS1, SSS2). Dashed line: boundary between Southern Brazilian Bight (SSB) and Southern Subtropical Shelf (SSS).

For the identification of organisms and their classification regarding to distribution in the marine environment of the Brazilian coast, the works of Cupp (1943), Desikachary (1959), Balech (1988); Balech *et al.* (1984), Round *et al.* (1990), Silva-Cunha & Eskinazi-Leça (1990), Torgan & Biancamano (1991), Hasle & Syvertsen (1997), Tomas (1997), Cardoso (1998) and Faust & Gullidge (2002) were used for the phytoplankton, and those of Rose (1933), Boltovskoy (1981, 1999), Conway *et al.* (2003), Johnson & Allen

(2005) and Conway (2012a, 2012b) for the protozooplankton and zooplankton.

Statistical analysis

The analysis of plankton data followed the procedure described by Field *et al.* (1982) and Lindley & Williams (1994), which used a 10% higher rate of occurrence of taxa per sample. The Primer® program was used to compose a 2-dimensional scatter plot from the matrices of Bray-Curtis coefficients, were calcu-

lated for similarities between the taxa occurring within samples and the distributions of taxa between samples for each tow, by MDS (non-metric multidimensional scaling). PERMANOVA (One-way) analysis was used to test the null hypothesis that the abundance and species richness levels of a species/taxa did not differ between sample groups, and Simper analysis was used to compare the main taxa occurring within samples, and the distributions between samples for each tow. PERMANOVA data were interpreted as indicating areas with significant different plankton groupings, and the composition of different groupings. Multivariate correlation analysis (Statistic®) was used and negative/positive correlation trophic plankton community.

RESULTS

A gradient in surface temperature of the water was observed, with higher temperatures to the north (SSB1 - 23.6 to 24.5°C), intermediate temperatures in the central region (SSB2, SSB3 and SSB4 - 21.2 to 23.4°C) and lower temperatures to the south (SSS1 and SSS2 - 19.4 to 21.3°C). We have identified 75 phytoplankton taxa, 26 zooplankton taxa, 10 protozooplankton taxa and fish eggs. The size of plankton is also a limitation sampling by VOR, because the size mesh and pico and nanoplankton rarely were collected, so, the community of plankton predominant is microphytoplankton (dinoflagellates, diatoms, cyanobacteria), copepods (zooplankton) and tintinnids (Ciliophora) (Table 2). The major percentage from VOR survey of phytoplankton (48%) is from coastal waters, zooplankton (48.6%) and protozooplankton (60%) from the mixture of neritic/oceanic waters (Figs. 2-3). The result had shown that there is bottom up effect between phytoplankton and zooplankton and both of them densities tended to decrease in a north-south region (Fig. 4). For tows involving the coast and the part of ocean (transects SSB3 and SSB4), the zooplankton showed a tendency to increase in density towards the ocean. All the tows made in north-south direction were more extensive than in the east-west direction, providing greater variations in spatial distribution of the plankton.

The planktonic community showed high similarity among transects, with representation of groupings between SSB1 and SSB2, SSS1 and SSS2 and an isolated group in SSB4 (MDS coefficient of stress: 0.18) (Fig. 5). Significant differences were also observed between the transect in the north region (SSB), and higher similarity in transects located in the south region (SSS) (Pseudo-F: 5.176) (Table 3). The taxa with more significant abundance among the transect analysed are shown in Table 4. The difference

for transect SSB1 in relation to the other transects is due to the high densities and predominance of the nanoplanktonic cyanobacteria *Johannesbaptistia* sp. This group was also important in transect SSB2, but with greater representation of other species (*Proboscia alata*, *Dictyocha fibula*, *Neoceratium fuscus*, *Thalassionema bacillare*) due the position of this transect, the farthest from the coast. Transect SSB3 presented high density of cyanobacteria at the beginning of the tow (*Thrichodesmium erythraeum*, *T. thiebautii*), followed by high density of diatoms (*Hemiaulus hauckii*) and dinoflagellates (*Prorocentrum sigmoides*), whereas that of transect SSB4 *Thalassiosira* spp. showed high density throughout the route. Transects SSB3 and SSB4 were similar due to their intermediate location, and because the tow direction was from the coast towards the ocean. Despite these differences, the distribution of phytoplankton was homogeneous for the transects located in SSB. For SSS the silicoflagellate *Octotactis octonaria* was predominant in SSS1 and SSS2.

For the protozooplankton, the predominant taxa in the southeastern region (SBB) were tintinnids (*Salpingella* sp.), which contributed to the high densities in SSB1, and zooplankton such as copepods there is high densities, as well as the presence of representatives of other groups such as Cladocera, Chaetognatha and Tunicata. In SSB2, copepods (*Paracalanus* sp., *Temora turbinata*, *Oithona* sp. and *Oncaea* sp.) and protozooplankton (*Codonellopsis*, *Salpingella* and *Dictyocysta*) had similar densities. Copepods (*Temora* and *Oncaea*) and Cladocera (*Penilia avirostris*) were the only groups observed in SSB3, and were low in number. In SSB4, copepods were predominant near the coast, while Appendicularia, Chaetognatha and Cladocera were representative in waters farther away from the coast. For the SSS region, Copepoda were the predominant group, represented by *Oithona*, *Oncaea*, *Clausocalanus* and *Centropages velificatus* in both transects of SSS1 and SSS2. Other groups of note were the Protozoa *Codonellopsis*, euphausiids, hyperiids and decapod larvae.

A positive correlation was observed between small-sized herbivorous zooplankton or those with microphagous habits (*Penilia avirostris*, *Salpingella* sp.) and the coastal phytoplankton (first axis). For other groups, such as amphipods hyperiids, euphausiids, and the copepods *Oithona* sp. and *Clausocalanus* sp., the distribution was associated with the habitats oceanic and oceanic/neritic (Fig. 6) and not with the trophic relationships.

DISCUSSION

The values for sea surface temperature showed differences between the studied regions, with a remarkable

Table 2. Classification of phytoplankton, protozooplankton and zooplankton observed in SSB (Southern Brazilian Bight) and SSS (Southern Subtropical Shelf).

Heterokontophyta	Dinophyta	Copepoda
<i>Coscinodiscophyceae</i>	<i>Ornithocercus magnificus</i>	<i>Acartia</i> sp.
<i>Thalassiosira</i> spp.	<i>Amphisolenia bidentata</i>	<i>Calanoides</i>
<i>Skeletonema</i> sp.	<i>Neoceratium fusus</i>	<i>Calanus</i> sp.
<i>Cerataulina pelágica</i>	<i>Neoceratium lineatum</i>	<i>Centropages velificatus</i>
<i>Cerataulina dentata</i>	<i>Neoceratium macroceros</i>	<i>Clausocalanus</i> sp.
<i>Corethron criophilum</i>	<i>Neoceratium hexacanthum</i>	Copepoda sp.
<i>Dactyliosolen</i> cf. <i>blavyanus</i>	<i>Neoceratium pentagonum</i>	Copepoda (eggs)
<i>Pseudosolenia calcar avis</i>	<i>Neoceratium horridum</i>	Copepoda (nauplii)
<i>Proboscia alata</i>	<i>Neoceratium furca</i>	Copepoda (juveniles)
<i>Rhizosolenia imbricata</i>	<i>Neoceratium condelabrum</i>	<i>Labidocera</i> sp.
<i>Rhizosolenia setigera</i>	<i>Neoceratium trichoceros</i>	<i>Paracalanus</i> sp.
<i>Rhizosolenia hebetata</i>	<i>Neoceratium</i> sp.	<i>Pseudocalanus</i> sp.
<i>Rhizosolenia</i> sp.	<i>Neoceratium tripos</i>	<i>Undinula</i> cf. <i>vulgaris</i>
<i>Bacteriastrum hyalinum</i>	<i>Alexandrium fraterculus</i>	Calanidae
<i>Bacteriastrum delicatulum</i>	<i>Scrippsiella trochoidea</i>	<i>Temora turbinata</i>
<i>Chaetoceros aequatorialis</i>	<i>Podolampas palmipes</i>	<i>Temora stylifera</i>
<i>Chaetoceros</i> cf. <i>brevis</i>	<i>Protopteridinium claudicans</i>	Poecilostomatoida
<i>Chaetoceros decipiens</i>	<i>Protopteridinium elegans</i>	<i>Corycaeus</i> sp.
<i>Chaetoceros diversus</i>	<i>Prorocentrum</i> cf. <i>mexicanum</i>	Harpacticoida
<i>Chaetoceros peruvianus</i>	<i>Prorocentrum micans</i>	<i>Oithona oswaldocruzi</i>
<i>Chaetoceros subtilis</i>	<i>Prorocentrum gracile</i>	<i>Oithona</i> sp.
<i>Chaetoceros whigamli</i>	<i>Prorocentrum minimus</i>	<i>Oithona plumifera</i>
<i>Chaetoceros eibenii</i>	<i>Pyrocystis obtusa</i>	<i>Oncaea</i> sp.
<i>Leptocylindrus danicus</i>	<i>Pyrocystis lunula</i>	<i>Oncaea media</i>
<i>Ditylum brighterii</i>	Chromophyta	<i>Oncaea mediterranea</i>
<i>Hemiaulus hauckii</i>	Dictyochophyceae	<i>Microsetella</i> sp.
Fragilariophyceae	<i>Dictyocha fibula</i>	Branchipoda-Cladocera
<i>Lioloma pacificum</i>	<i>Dictyocha</i> sp.	<i>Penilia avirostris</i>
<i>Thalassionema nitzschioides</i>	<i>Octotactis octonaria</i>	<i>Podon</i> sp.
<i>Thalassionema bacillare</i>	Cyanophyta	Tunicata
<i>Climacospheonia</i> sp.	<i>Synechococcus</i> sp.	<i>Appendicularia</i> sp.
Bacillariophyceae	<i>Johannesbaptistia</i> sp.	Chaetognatha
Naviculaceae	<i>Richelia intracellularis</i>	Tecossoma
<i>Pleurosigma aestuarii</i>	<i>Katagnymene pelágica</i>	Euphausidae
<i>Cylindroteca closterium</i>	<i>Trichodesmium erythraeum</i>	Hyperidae
<i>Nitzschia longissima</i>	<i>Trichodesmium hildebrandtii</i>	Decapoda
<i>Pseudo-nitzschia</i> - complex <i>delicatissima</i>	<i>Trichodesmium</i> sp.	Fish eggs
<i>Pseudo-nitzschia</i> - complex <i>seriata</i>	<i>Oscillatoria</i> sp.	
<i>Synedra</i> sp.	Oscillatoriales 1	
<i>Entomoneis</i> sp.		
Protozooplankton	<i>Dictyocysta</i> sp.	<i>Salpingella</i> cf. <i>acuminatoides</i>
<i>Codonellopsis</i> sp.	<i>Dadayiella ganymedes</i>	<i>Tintinopsis campanula</i>
<i>Codonella</i>	<i>Eutintinus</i> sp.	<i>Xystonellopsis gaussi</i>
<i>Codonellopsis schabi</i>	<i>Salpingella</i> sp.	

gradient from north to south. The transects of region SSB (of SSB1 SSB4) was characterized by coastal and warm continental shelf waters, or the influence of the Brazil Current, due to higher temperatures, in according to Silveira *et al.* (2000). In the south region (SSS1 and SSS2), the influence of coastal water, with the La Plata front (Piola *et al.*, 2000) may have contributed to lower

temperature observed. This thermal gradient was probably a primary factor forcing the separation of the groups of transects between the regions SSB and SSS observed in the ordering analyses (MDS), differentiating the densities and groups of phytoplankton and zooplankton.




Figure 2. Classification of phytoplankton (taxonomy) according to their habitat distributions (oceanic, oceanic/neritic, coastal waters and coastal water/estuarine) and its occurrence in SSB (Southern Brazilian Bight), SSB/SSS (both region), SSS (Southern Subtropical Shelf).




Figure 3. Classification of zooplankton, protozooplankton and other groups (taxonomy) according to habitat distributions (oceanic, oceanic/neritic, neritic, neritic/coastal waters and coastal water/estuarine) and its occurrence in SSB (Southern Brazilian Bight), SSB/SSS (both regions), SSS (Southern Subtropical Shelf).




Figure 4. Total phytoplankton (log10) and zooplankton (log+1) density each 5 nm (9.26 km) between towing areas (transects: SSB1, SSB2, SSB3, SSB4, SSS1, SSS2).

For the Brazilian coastal zone, these are the first results related to the category of continuous plankton sampling with the use of VOR and microplankton was the fraction with the highest yields in the VOR tows and was also common in the North Sea and North Atlantic Ocean (Lindley & Williams, 1994; Beaugrand *et al.*, 2000; Reid *et al.*, 2003; Richardson *et al.*, 2006). Besides the presence of diatoms and dinoflagellates in transects SSB1, SSB2 and SSB3, the nanophytoplankton (*Johannesbaptistia* sp.) was observed, accompanied by *Thrichodesmium erythraeum*, *Thrichodesmium thiebautii*, which were prevalent in high density. The proximity of transects SSB1 and SSB2 to the coast may favour the occurrence of *Johannesbaptistia* sp., due to its association with an environment with higher loads of organic matter (Cohen *et al.*, 1986), which is also commonly observed in tropical environments upon presence of river inflows (Bauer *et al.*, 2008). The genus *Thrichodesmium* characterizes high productivity in the oceanic region (Gallon *et al.*, 1996) and occurs from the northeast of Brazil (Koenig

& Macedo, 1999) to the southeastern and southern (Brandini & Fernandes, 1996), but with low density in Subantarctic Water (Brandini, 1988).

For the transects of the arrangement farther to the south (SSB4; SSS1 and SSS2), there was observed the high densities of diatoms and silicoflagellates, which are related to a higher concentration of silicate in this region, especially during the winter, with the two largest river inputs in the south region (Río de la Plata and the Lagoa dos Patos Estuary (Piola *et al.*, 2000), in addition to the high concentration of nutrients (Rigual-Hernandez *et al.*, 2010). These phytoplanktonic groups have already been observed in the same period, associated with the cold water of Subantarctic origin (Brandini, 1998). As a result, the high prevalence of these groups is reflected in the subsequent trophic levels, as they represent the main diet of copepods common to the Brazilian coast (Lopes, 1998).

The vast majority of zooplanktonic carcasses observed in all transects were those of the copepods group (<2 mm) (*Acartia* sp., *Calanus* sp., *Centropages*




Figure 5. MDS analysis (Bray Curtis similarity - log+1; Stress: 0.18) to transects in base of phytoplankton and zooplankton densities (frequency of >10%). Sections of 5 nm to each transect (SSB1, SSB2, SSB3, SSB4, SSS1, SSS2).

Table 3. Plankton transects PERMANOVA analysis (P (perm), *non-significant difference).

P (perm)	SSB1	SSB2	SSB3	SSB4	SSS1	SSS2
SSB1	-					
SSB2	0.019	-				
SSB3	0.006	0.005	-			
SSB4	0.002	0.001	0.021	-		
SSS1	0.001	0.001	0.001	0.002	-	
SSS2	0.007	0.003	0.019	0.017	0.071 *	-
Perms: 998	Pseudo-F: 5,176	P (perm): 0.001		P(MC): 0.001		

velificatus, *Clausocalanus* sp., *Paracalanus* sp., *Pseudocalanus* sp., *Temora turbinata*, *Undinula vulgaris*, *Microsetella* sp., *Oithona* sp., *Oithona plumifera*, *Oncaea* sp., *Oncaea mediterranea*) of the shelf. The small size of dominant species is related to the Coastal Water (CW) and the mixing with Tropical Water (TW), and some species are restricted to estuaries and coastal circulation, their distribution being limited under the influence of oceanic waters (Lopes *et al.*, 2006). The greater abundance of copepods in the south region was also reported by Nunes & Resgalla Jr. (2012). Cladocera was not observed in the samples of the VOR that it only operates in a narrow depth range, which is outside the

preferential distribution of the group even though Resgalla Jr. & Montú (1993) and Muxagata & Montú (1999) reported in the southern coast and southeastern Brazilian. This fact may be related to the limitation of the VOR that it only operates in a narrow depth range, which is outside the preferential distribution of the group.

The prevalence of Chaetognatha (SSB1, SSB2, SSB4, SSS1), Copepoda (*Acartia* sp., *Temora stylifera*, *Temora turbinata*, *Oithona plumifera*, *Oithona* sp., *Paracalanus* sp.) in SSB1 and Cladocera (*Penilia avirostris*) (SSB1, SSB2 and SSB3), are indicative of Coastal Water (WC) and its association with estuarine regions (Resgalla Jr. & Montú, 1993; Resgalla Jr., 2011). The genera *Paracalanus* and *Temora*, present in SSB1, are consumers of the community of phytoplankton and microzooplankton in regions close to the coastal upwelling zone (Lopes *et al.*, 1998). Also noteworthy is the high abundance of protozooplankton, especially Ciliophora (*Codonellopsis schabi*, *Salpingella cf. acuminatoides*), in transects of the north region, which is the region closest to the Cabo Frio upwelling, in the State of Rio de Janeiro. This group is dominated by herbivorous, with nanoplankton being an important item in its diet (Gowing & Garrison, 1992).

Table 4. The percentage contribution (%) of abundance of plankton by Simper analysis between transects SSB1, SSB2, SSB3, SSB4, SSS1 and SSS2.

Rate/Transects	SSB1	SSB2	SSB3	SSB4	SSS1	SSS2
Average similarity (%)	57.80	33.81	38.45	30.74	45.74	40.61
% Similarity each taxa						
<i>Johannesbaptistia</i> sp.	14.52	10.60				
<i>Dictyocha fibula</i>	11.67	14.14				
<i>Proboscia alata</i>	8.62	14.75				
<i>Neoceratium fusus</i>	7.84	13.02				
<i>Chaetoceros decipiens</i>	7.77					
<i>Thalassionema bacillare</i>		11.14		27.52		
<i>Hemiaulus hauckii</i>			38.14			
<i>Prorocentrum sigmoides</i>			23.24			
<i>Corethron criophilum</i>			10.07			
<i>Thalassiosira</i> spp.				55.52	18.39	
<i>Rhizosolenia setigera</i>					34.49	
<i>Octotactis octonaria</i>					19.12	29.93
<i>Thalassionema nitzschioides</i>						16.53
<i>Rhizosolenia setigera</i>						12.17
<i>Appendicularia</i> sp.		4.27				
<i>Penilia avirostris</i>		4.18				
<i>Oithona</i> sp.		2.77				




Figure 6. Multivariate correlation between phytoplankton and zooplankton density (>10%, transf. data: log).

The efficiency of Continuous Plankton Sampling has been compared with that of traditional samplers in different studies, and has proven to be robust (Batten *et al.*, 2003b; Hunt & Hoise, 2003). In different studies of plankton involving the CPR, the diversity and abundance have presented high confidence in time series analyses for monitoring climate change in the North Atlantic (Edwards *et al.*, 2010; Mcleod *et al.*, 2012). In addition, plankton is currently the only group to have been monitored continuously since the 1940s, and multi-decadal analyses have already been established, due to the quantity and robustness of the data collected by this method. In according to the data provided by the VOR, on the continental shelf of Southeastern Brazil, the planktonic community was dominated by organisms smaller than 2 mm, whose distribution has reached the South region. Estuarine and coastal waters phytoplanktonic species showed a predominance of cyanobacteria and diatoms in SSB, and silicoflagellates in SSS, while the distribution of zooplankton was characterized by small copepods in SSB and genres of larger size in SSS, but all with neritic/oceanic characteristics.

ACKNOWLEDGMENTS

To the Captains of the fishing boats the Mtanos Seif and Nicassio da Costa and the Itajaí Fishermens' Union (SINDIPI), who enabled the tows, to scientific observers Matheus Bella and Fernão Weiss, SAHFOS team (Plymouth, UK) where the VOR analyses were performed. To profesor Tito C. Marques for his assistance in the statistical analyses, CAPES (Doctorate student, Sandwich course, process no.: 99999.003694/2014-03). To the National Council for Scientific and Technological Development (CNPq) for the research grant, process no. 305795/2014-7. The cruises were funded by IGEPESCA (Edital Ciências do Mar/CAPES-2009).

REFERENCES

- Alvariño, A. 1980. El plancton del Atlántico suroeste, dinámica y ecología. Bolm. Inst. Oceanogr. São Paulo, 29: 15-26.
- Araújo, H.M.P. & V.A. Ribeiro. 2005. Distribuição das espécies de Chaetognatha na plataforma continental de Sergipe e Alagoas. Braz. J. Aquat. Sci. Technol., 9: 19-23.
- Balech, E., R. Akselman, H.R. Benavides & R.M. Negri. 1984. Suplemento a los dinoflagelados del Atlántico Sudoccidental. INIDEP, Mar del Plata, 20 pp.
- Balech, E. 1988. Los dinoflagelados del Atlántico sudoccidental. Publ. Espec., Inst. Español Oceanogr., 310 pp.
- Batten, S.D., A.W. Walne, M. Edwards, S.B. Groom. 2003a. Phytoplankton biomass from continuous plankton recorder data: an assessment of the phytoplankton colour index. J. Plankton Res., 25: 697-702.
- Batten, S.D., R. Clark, J. Flinkman, G. Hays, E. John, A.W.G. John, T. Jonas, J.A. Lindley, D.P. Stevens & A. Walne. 2003b. CPR sampling: the technical background, materials and methods, consistency and comparability. Prog. Oceanogr., 58: 193-215.
- Beaugrand, G., F. Ibanez & P.C. Reid. 2000. Spatial, seasonal and long-term fluctuations of plankton in relation to hydroclimatic features in the English Channel, Celtic Sea and Bay of Biscay. Mar. Ecol. Prog. Ser., 200: 93-102.
- Boltovskoy, D. 1981. Atlas del zooplancton del Atlántico Sudoccidental y métodos de trabajo con el zooplancton marino. INIDEP, Mar del Plata, 936 pp.
- Boltovskoy, D. 1999. South Atlantic zooplankton. Backhuys Publishers, Leiden, 1705 pp.
- Brandini, F.P. 1988. Composição e distribuição do fitoplâncton na Região Sueste e suas relações com as massas d'água (Operação Sueste I - Invemo/1982). Ciênc. Cult., 40: 337-341.
- Brandini, F.P. 1990. Hydrography and characteristics of the phytoplankton in shelf and oceanic waters off southeastern Brazil during winter (July/August-1982) and summer (February/March-1984). Hydrobiologia, 196: 111-148.
- Brandini, F.P. & L.F. Fernandes. 1996. Microalgae of the continental shelf off Paraná State, southeastern Brazil: a review of studies. Rev. Bras. Oceanogr., 44: 69-80.
- Cardoso, L.S. 1998. Dinoflagelados da Ilha do Arvoredo e da Praia de Ponta das Canas, Santa Catarina, Brasil. Biociências, 6: 3-54.
- Colebrook, J.M. 1960. Continuous plankton records: methods of analysis, 1950-1959. Bull. Mar. Ecol., 41: 51-64.
- Cohen, Y., B.B. Jorgensen, N.P. Revsbech & R. Poplawski. 1986. Adaptation to hydrogen sulfide of oxygenic and anoxygenic photosynthesis among Cyanobacteria. Appl. Envir. Microbiol., 51: 398-407.
- Conway, D.V.P. 2012a. Identification of the copepodite developmental stages of twenty-six North Atlantic copepods. Occasional Publication Mar. Biol. Assoc. UK, 21: 35 pp.
- Conway, D.V.P. 2012b. Marine zooplankton of southern Britain. Part 1. Radiolaria, Heliozoa, Foraminifera, Ciliophora, Cnidaria, Ctenophora, Platyhelminthes, Nemertea, Rotifera and Mollusca. A.W.G. John (ed.).

- Occasional Publication of the Mar. Biol. Assoc. U.K., 25: 138 pp.
- Conway, D.V.P., R.G. White, J. Hugues-Dit-Ciles, C.P. Gallienne & D.B. Robins. 2003. Guide to the coastal and surface zooplankton of the south-western Indian Ocean. Occasional Publication of the Mar. Biol. Assoc. U.K., 15: 356 pp.
- Cupp, E.E. 1943. Marine plankton diatoms of the west coast of North America. University of California Press, Berkeley and Los Angeles, 235 pp.
- Domingos-Nunes, R. & C. Resgalla Jr. 2012. The zooplankton of Santa Catarina continental shelf in southern Brazil with emphasis on Copepoda and Cladocera and their relationship with physical coastal processes. *Lat. Am. J. Aquat. Res.*, 4: 893-913.
- Edwards M. & A.J. Richardson. 2004. Impact of climate change on marine pelagic phenology and trophic mismatch. *Nature*, 430.
- Edwards, M., P.C. Reid & B. Planque, 2001. Long-term and regional variability of phytoplankton biomass in the Northeast Atlantic (1960-1995). *ICES J. Mar. Sci.*, 58: 39-49.
- Edwards, M., G. Beaugrand, G.C. Hays, J.A. Koslow & A.J. Richardson. 2010. Multi-decadal oceanic ecological datasets and their application in marine policy and management. *Trends Ecol. Evol.*, 25: 602-610.
- Faccin, J., D.M.C. Gama Cunha, R. Barddal & C. Resgalla Jr. 2014. Development of an oceanographic towing vehicle adapted for fishing craft: Prototype and protocol for use. *Meth. Oceanogr.*, 9: 61-74.
- Faust, M.A. & R.A. Gulledge. 2002. Identifying harmful marine dinoflagellates. *Natl. Mus. Nat. Hist. Wash.*, 144 pp.
- Fernandes, L.F. 1999. Tintininos (Ciliophora - Subordem Tintinnina) de águas subantárticas e antárticas entre a Argentina e a Península Antártica (35°S-62°S) (November de 1992). *Rev. Bras. Oceanogr.*, 47: 155-171.
- Fernandes, L.F. 2004. Tintininos (Ciliophora, Tintinnina) de águas subtropicais na região Sueste-Sul do Brasil. I. Famílias Codonellidae, Codonellopsidae, Coxliellidae, Cyttarocylidae, Epiplocylidae, Petalotrichidae, Ptychocylidae, Tintinnididae e Undellidae. *Rev. Bras. Zool.*, 21: 551-576.
- Fernandes, L.F. & F.P. Brandini. 2004. Diatom associations in shelf waters off paraná state, southern brazil: annual variation in relation to environmental factors. *Braz. J. Oceanogr.*, 52: 19-34.
- Field, J.G., K.R. Clarke & P. Mayzaud. 1982. A practical strategy for analysing multispecies distribution patterns. *Mar. Ecol. Prog. Ser.*, 8: 37-52.
- Gallon, J.R., D.A. Jones & T.S. Page. 1996. *Trichodesmium*, the paradoxical diazotroph. *Algological Stud.*, 83: 215-243.
- Gomes, N.A., E. Eskinazi-Leça & M.G.G. Silva-Cunha. 1989. Ocorrência de *Auriculia complexa* (Bacillariophyceae) na plataforma continental de Pernambuco. *Acta Bot. Bras.*, 3(2): 7 pp.
- Head, E.J.H. & P. Pepin. 2010a. Monitoring changes in phytoplankton abundance and composition in the Northwest Atlantic: a comparison of results obtained by continuous plankton recorder sampling and colour satellite imagery. *J. Plankton Res.*, 12: 1649-1660.
- Head, E.J.H. & P. Pepin. 2010b. Spatial and inter-decadal variability in plankton abundance and composition in the Northwest Atlantic (1958-2006). *J. Plankton Res.*, 12: 1633-1648.
- Hosie, G.W., M. Fukuchi & S. Kawaguchi. 2003. Development of the Southern Ocean Continuous Plankton Recorder survey. *Prog. Oceanogr.*, 58: 263-283.
- Hunt, B.P.V. & G.W. Hosie. 2003. The Continuous Plankton Recorder in the Southern Ocean: a comparative analysis of zooplankton communities sampled by the CPR and vertical net hauls along 140°E. *J. Plankton Res.*, 12: 1561-1579.
- John, E.H., S.D. Batten, D. Stevens, A.W. Walne, T. Jonas & G.C. Hays. 2002. Continuous plankton records stand the test of time: evaluation of flow rates, clogging and the continuity of the CPR time-series. *J. Plankton Res.*, 9: 941-946.
- Johnson, W.S. & D.M. Allen. 2005. Zooplankton of the Atlantic and Gulf coasts: a guide to their identification and ecology. Johns Hopkins University Press, New York, 369 pp.
- Klein, V.L.M. 1977. Sobre a composição e abundância relativa do plâncton, na plataforma continental do estado do Ceará. *Arq. Ciênc.Mar*, 17: 21-27.
- Koenig, M.L. & S.J. Macedo. 1999. Hydrology and phytoplankton community structure at Itamaracá-Pernambuco (Northeast Brazil). *Braz. Arch. Biol. Technol.*, 2: 381-392.
- Koenig, M.L., B.E.B. Wanderley & S.J. Macedo. 2009. Microphytoplankton structure from the neritic and oceanic regions of Pernambuco State - Brazil. *Braz. J. Biol.*, 69: 1037-1046.
- Lindley, J.A. & S.D. Batten. 2002. Long-term variability in the diversity of North Sea zooplankton. *J. Mar. Biol. Assoc., U.K.*, 82: 31-40.
- Lindley, J.A. & R. Williams. 1994. Relating plankton assemblages to environmental variables using instruments towed by ships-of-opportunity. *Mar. Ecol. Prog. Ser.*, 107: 245-262.
- Lopes, R.M., R. Vale & F.P. Brandini. 1998. Composição, abundância e distribuição espacial do zooplâncton no complexo estuarino de Paranaguá durante o inverno de 1993 e o verão de 1994. *Rev. Bras. Oceanogr.*, 46: 195-211.

- Lopes, R. M., M. Katsuragawa, J.F. Dias, M.A. Montú, J.H. Muelbert, C. Gorri & F.P. Brandini. 2006. Zooplankton and ichthyoplankton distribution on the southern Brazilian shelf: an overview. *Sci. Mar.*, 70: 189-202.
- Lucas, C.E. 1941. Continuous Plankton Records: phytoplankton in the North Sea 1938-39. Part. I Diatoms. *Hull Bull. Mar. Ecol.*, 8: 19-46.
- McLeod, D.J., G.M. Hallegraeff, G.W. Hosie & A.J. Richardson. 2012. Climate-driven range expansion of the red-tide dinoflagellate *Noctiluca scintillans* into the Southern Ocean. *J. Plankton Res.*, 34: 332-337.
- Moser, G.A.O. & S.M.F. Gíanesella-Galvão. 1997. Biological and oceanographic upwelling indicators at Cabo Frio (RJ). *Rev. Bras. Oceanogr.*, 45: 11-23.
- Muxagata, E. & M.A. Montú. 1999. Os Cladorera da plataforma continental sudeste brasileira: distribuição, densidade e biomassa (inverno de 1995). *Nauplius*, 7: 151-172.
- Olguin, H.F., D. Boltovskoy, C.B. Lange & F. Brandini, 2006. Distribution of spring phytoplankton (mainly diatoms) in the upper 50 m of the Southwestern Atlantic Ocean (30-61'S). *J. Plankton Res.*, 28: 1107-1128.
- Palma, E.D., R.P. Matano & A.R. Piola. 2008. A numerical study of the Southwestern Atlantic Shelf circulation: stratified ocean response to local and offshore forcing. *J. Geophys. Res.*, 113: C11010.
- Piola, A., E.D. Campos, O.O. Moller Jr., M. Charo & C. Martinez. 2000. Subtropical shelf front off eastern South America. *J. Geophys. Res.*, 105: 6565-6578.
- Reid, P.C., J.M. Colebrook, J.B.L. Matthews & J. Aiken. 2003. The Continuous Plankton Recorder: concepts and history, from Plankton Indicator to undulating recorders. *Prog. Oceanogr.*, 58(2-4): 117-173.
- Resgalla Jr., C. 2011. The holoplankton of the Santa Catarina coast, southern Brazil. *Anais Acad. Bras. Ciênc.*, 83: 575-588.
- Resgalla Jr., C. & M. Montú. 1993. Cladoceros marinhos da plataforma continental do Rio Grande do Sul - Brasil. *Nauplius*, 1: 63-79.
- Resgalla Jr., C., C. De La Rocha, & M. Montú. 2001. The influence of Ekman transport on zooplankton biomass variability off southern Brazil. *J. Plankton Res.*, 23: 641-650.
- Richardson, A.J., A.W. Walne, A.W.G. John, T.D. Jonas, J.A. Lindley, D.W. Sims, D. Stevens & M. Witt. 2006. Using continuous plankton recorder data. *Prog. Oceanogr.*, 68: 27-74.
- Rigual-Hernandez, A.S., M.A. Barcena, F.J. Sierro, J.A. Flores, I. Hernandez-Almeida, A. Sanchez-Vidal, A. Palangues & S. Heussner. 2010. Seasonal to inter-annual variability and geographic distribution of the silicoflagellate fluxes in the Western Mediterranean. *Mar. Micropaleontol.*, 77: 46-57.
- Rose, M. 1933. Faune de France. Copépodes pélagiques. *Fédération Française des Sociétés de Sciences Naturelles Office Central de Faunistique*, 26: 337 pp.
- Silveira, I.C.A., A.C.K. Schmidt, E.J.D. Campos, S.S. Godoi & Y. Ikeda. 2000. A Corrente do Brasil ao largo da costa leste brasileira (The Brazil Current off the Eastern Brazilian Coast). *Rev. Bras. Oceanogr.*, 48: 171-183.
- Souza, E.B., V.B. Costa, L.C.C. Pereira & R.M Costa. 2008. Microplankton de águas costeiras amazônicas: Ilha Canela (Bragança, PA, Brasil). *Acta Bot. Bras.*, 22: 626-636.
- Tomas, C.R. (ed.). 1997. Identifying marine phytoplankton. Academic Press, California, 857 pp.
- Villac, M.C. & D.R. Tenenbaum. 2010. The phytoplankton of Guanabara Bay, Brazil. I. Historical account of its biodiversity. *Biot. Neotrop.*, 10: 271-293.
- Warner, A.J. & G.C. Hays. 1994. Sampling by the continuous plankton recorder survey. *Prog. Oceanogr.*, 34: 237-256.

Received: 12 October 2015; Accepted: 12 June 2016