



Research Article

An update on female sizes and body condition of nesting olive ridley turtles (*Lepidochelys olivacea*) in La Escobilla Beach Sanctuary, Mexico

Alejandra Buenrostro Silva^{1,2} , Petra Sánchez Nava³ 

Jesús García-Grajales²  & María de Lourdes Ruiz Gómez³ 

¹Programa de Doctorado en Ciencias Agropecuarias y Recursos Naturales

Universidad Autónoma del Estado de México, Toluca, Estado de México, México

²Universidad del Mar Campus Puerto Escondido, San Pedro Mixtepec, Oaxaca, México

³Facultad de Ciencias, Universidad Autónoma del Estado de México

Toluca, Estado de México, México

Corresponding author: Petra Sánchez Nava (psn@uaemex.mx)

ABSTRACT. Body condition (BC) can affect the reproductive output and hatchling survival of nesting olive ridley turtles (*Lepidochelys olivacea*). However, little is known about the BC of nesting females during the arrival event at nesting sites. Therefore, this study aimed to evaluate the BC of females nesting in La Escobilla Beach Sanctuary (La Escobilla) during the 2021 nesting season. Nesting females were measured and weighed. The mean size for nesting *L. olivacea* was 60.41 ± 2.98 cm for straight carapace length, 52.69 ± 2.57 cm for straight carapace width, and 34.35 ± 3.94 kg for weight ($n = 483$). The mean (\pm standard deviation, SD) Fulton's/Bjorndal BC index estimated for nesting females was 1.54 ± 0.15 , while the mean (\pm SD) of body mass index was 2.18 ± 0.23 . Based on these indexes, 28 individuals were classified as in bad condition, 315 in good condition, and 19 in excellent condition. This first estimation of the olive ridley turtle body condition in a nesting colony showed that BC is a good body condition for most nesting females in La Escobilla. This study is a valuable baseline for nesting *L. olivacea's* health condition.

Keywords: *Lepidochelys olivacea*; arribada; health assessment; body mass; length; reproductive output

INTRODUCTION

The olive ridley turtle *Lepidochelys olivacea* is a pantropical marine species distributed worldwide in tropical and subtropical oceanic regions (Abreu-Grobois & Plotkin 2008). This species is currently listed as "Vulnerable" according to the International Union for Conservation of Nature Red List (Abreu-Grobois & Plotkin 2008) and classified as "Endangered" by Mexican laws (DOF 2010). Like most sea turtles, the olive ridley turtle has been severely affected by human activities, such as direct take and illegal trade and harvesting of eggs for local consumption, both considered to be one of the species's major threats

(Fonseca et al. 2009, Valverde et al. 2012, Bézy et al. 2016, Pheasey et al. 2020).

This species exhibits two nesting strategies, one associated with nesting in a solitary fashion and the other, a unique nesting behavior where hundreds to thousands of turtles emerge on nesting beaches in a synchronous event called "arribadas" (Cornelius et al. 1991, Bernardon & Plotkin 2007). In the Americas, the most notable arribadas nesting beaches are in Costa Rica (Cornelius et al. 1991, Plotkin et al. 1997, Campbell 1998, Hornarvar et al. 2008, Fonseca et al. 2009), Nicaragua (Steward 2001, Hope 2002), Panamá (Cornelius et al. 2007), and Mexico (Peñaflares et al. 2000, Campbell 2007).

Mexico has one of the most important nesting areas of the *L. olivacea* on La Escobilla Beach Sanctuary (La Escobilla) in Oaxaca, on the eastern Pacific coast (Márquez et al. 1996, Ocaña et al. 2012). This site was designated in 1986 as a protected natural area under the category of sanctuary to protect the olive ridley turtles. The area is under the management of the Comisión Nacional de Área Naturales Protegidas (CONANP, Márquez et al. 1996). Generally, the arribadas occur at a particular time of the year, between June and December, commonly once a month. Occasionally, an arribada can last a single day or up to 30 days (Ocana et al. 2012). In the eastern Pacific, the arribadas coincide annually with the rainy season (June-October, Cornelius 1986), as for La Escobilla.

Most sea turtle monitoring programs are focused on assessing population trends (Nel et al. 2013), or the protection of nesting sites (García et al. 2003, Gaona & Barragán 2016, Mazaris et al. 2017). However, there is little information on the health condition of the nesting females (e.g. individual size has decreased over time, Tuček et al. 2014). For example, in India, a declining trend in female sizes has been detected for olive ridleys (Shanker et al. 2003). Similar patterns had been shown for hawksbill turtles (*Eretmochelys imbricata*) in Mexico (Perez-Castaneda et al. 2007), loggerhead turtles (*Caretta caretta*) in Turkey (Ilgaz et al. 2007), and green turtles (*Chelonia mydas*) in Hawaii (Piacenza et al. 2016). While information shows that the olive ridley nesting population in Mexico is growing (Seminoff et al. 2008, Márquez et al. 2014), the condition of nesting females remains in a gap. Moreover, evaluation of female sizes and individual turtles' health is rare in Mexico despite their importance for improving conservation efforts (Nolte et al. 2020). Particularly, the knowledge of female body size will enable us to establish the relationship between reproductive output and body size (Hays & Speakman 1991, LeBlanc et al. 2014, LeGouvello et al. 2020a,b). Furthermore, body condition (BC) knowledge of nesting females is important because it can affect the reproductive output and hatchling's survival (Perrault et al. 2012); however, BC is considered a poor predictor to assess the adipose tissue (Kophamel et al. 2023).

BC is the physical status of an animal's body (Stevenson & Woods 2006). It is used to evaluate the fitness of animals in their environment (Taylor 1979, Peig & Green 2010). It also measures individual conditions, usually related to nutritional status and energy reserves (Harder & Kirkpatrick 1996 in Copeland 2004). Ecologists have used BC indexes to estimate an animal's nutritional state and a proxy for its

health condition (Ullman-Culleré & Foltz 1999). To estimate sea turtles' BC, morphometric data collection from many individuals is required (Thomson et al. 2009); hence, a BC index is a metric derived from the relationship between an animal's length and its mass (Labrada-Martagon et al. 2010). Until now, little is known about the BC of nesting females that emerge on nesting beaches in arribadas. Ramírez-Villanueva et al. (2023) documented the BC of a few females in the territorial sea off nesting beaches on the Pacific coast of Mexico off Oaxaca. Changes in the BC, resulting from changes in the biochemical composition of the tissue and mobilization of energy reserves, may be related to season, life-history traits, health status, or exposure to stressors (Barton et al. 2002). Hence, reproductive success, survival, and population dynamics depend on the BC of the organisms (Jones et al. 1999, Stevenson & Woods 2006).

Several approaches have been employed to estimate and evaluate individual or group BC in turtles (Stevenson & Woods 2006), including mass-length relationships (Bjørndal et al. 2000, Jessop et al. 2004), blood sample analysis (Stamper et al. 2005), examination of the plastron (Thomson et al. 2009), epibiotic barnacle loading (Deem et al. 2009, Flint et al. 2010, Nájera-Hillman et al. 2012), and adipose tissue estimation by bioelectrical impedance analysis (Kophamel et al. 2023). Moreover, these BC indexes have even been linked to life history traits such as mate selection, territorial extension, and mortality (Green 2001). The Fulton's condition factor is the most widely used BC index in sea turtle health-assessment studies (Ricker 1975, Bjørndal et al. 2000). This study emerges from fishery sciences and is based on the assumption that all parts of an ideal theoretical fish grow similarly (isometric growth) (LeCren 1951, Bjørndal et al. 2000). Thus, generates an index that can be related to the animal's length, dissociating both body size and BC and obtaining an abstract measurement of energy reserves (LeCren 1951, Cone 1989). Nonetheless, this assumption is rarely true (LeCren 1951). Healthy individuals are more likely to reproduce because they have high energy reserves than animals with low conditions (Schulte-Hostedde et al. 2005). Therefore, these individuals are essential for the population's health and supporting proper ecosystem functioning (Meffe 1999, Munson & Karesh 2002). This study aimed to evaluate the female sizes and BC of the nesting olive ridley turtle in La Escobilla and provide a baseline for understanding this population and future trends.

MATERIALS AND METHODS

Study area

La Escobilla Beach is within the municipality of Santa Maria Tonameca (Fig. 1) in the state of Oaxaca, on the southwest Mexican Pacific coast (15°43'37''N, 96°44'49''W). The climate is warm-subhumid, with a mean annual temperature of 28°C. The mean annual precipitation is 1000 mm, with rains falling between May and October (García 1981). The beach is approximately 25 km long, and the arribadas turtles nest along an 8-km strip at the beach's western end (Ocaña et al. 2012).

Data collection

This study was conducted from August through December 2021, during the olive ridley turtle nesting season on La Escobilla. Nesting females were approached at night approximately 5 min after nest-building behavior ceased and egg-laying activity had begun. If the female did not appear in an egg-laying trance at the time of the first approach, a 5-min wait was added to the egg-laying disturbance (Whiting et al. 2007). Each turtle was evaluated as prone and supine to determine that the selected female had no injuries, mutilations, or obvious signs of illness. One examiner held the animal to avoid excessive movement, and the other performed a detailed systematic inspection in a cranial-caudal dorsoventral orientation (Resendiz et al. 2018). All those organisms that presented evidence of fractures and injuries were excluded from this study.

Turtles were examined using two methodologies: 1) straight carapace length (SCL), straight carapace width (SCW), and thickness of carapace (TC) were measured using a 950 mm long Haglöf tree caliper (Model Mantax Blue; estimated measurement error ± 0.5 cm) (Le Gouvello et al. 2020a, Lamont & Johnson 2021). The SCL was measured from the nuchal notch to the most posterior portion of the rear marginals. The SCW was measured from the widest part of the carapace, and the TC was measured from the middle of the plastron to the highest part of the carapace; 2) the curved carapace length (CCL), curved carapace width (CCW), and diameter of body (circumference, Cir), in the same carapace locations for straight measures. All measurements were taken with a flexible tape (measurement error ± 0.5 cm; Le Gouvello et al. 2020a, Lamont & Johnson 2021). Finally, the weight (W) was taken in kilograms for each turtle by suspending the turtle by ropes and a portable stretcher attached to a digital scale (instrument error ± 0.5 kg), anchored to a metallic tripod (Lamont & Johnson 2021).

Body condition estimates

The BC of nesting females was estimated using three different BC index: 1) Fulton index (K), 2) BC index modified by Bjorndal et al. (2000) (Bjorndal index, hereafter), and 3) body mass index (BMI). Fulton's index has been estimated for several sea turtle species and is calculated from the length-weight relationship (Beverton & Holt 1957). Bjorndal et al. (2000) proposed an index to determine the BC for green turtles (*C. mydas*) in the Bahamas National Park system, which was then taken as a reference to determine BC. Additionally, we implemented a new approximation to the BMI, commonly used in veterinary sciences and cattle ranching (Salazar-Cuytún et al. 2020).

The K was calculated as:

$$K = \frac{W}{L^b} \times 10^n$$

where b is the scaling exponent (isometric and therefore equal to three), the result is multiplied by 10 and raised to n to achieve a unit.

The Bjorndal index was calculated as $BC = [\text{body mass } (W) \times \text{SCL}^{-3}] \times 10,000$, where body mass is the animal's W in kilograms, SCL is the straight carapace length, and 3 is the scaling exponent, which is isometric and therefore, equal to three.

The BMI was calculated as

$$BMI = [W \text{ (kg)} / TC \text{ (mm)}] / SCL \text{ (mm)} / 10$$

where W is the animal's W in kilograms, thickness of the carapace (TC) is the measure from the middle of the plastron to the highest part of the carapace in mm, and SCL is the straight carapace length. Traditionally, this formula is applied to measure BMI in sheep, horses, swine, and other farm animals (Salazar-Cuytún et al. 2020), using the height of the withers; here, we replaced this for TC.

The BC of nesting turtles, based on the three BC indexes, was characterized by Castro et al. (2001) as a) bad condition, individuals whose BC is less than the arithmetic mean of the sample minus a standard deviation; b) good condition, individuals whose BC is equal to or greater than the previous value and less than or equal to the sum of the mean plus a standard deviation; and c) excellent condition, representing individuals whose BC is greater than the sum of the average and a standard deviation.

Data analysis

Simple linear regression was used to analyze the relationship between W and each measurement variable (SCL, SCW, CCL, CCW, Cir). Mann-Whitney tests




Figure 1. Location of La Escobilla Beach Sanctuary, Oaxaca, Mexico.

were used to identify differences between BC estimates. As a product of applying the formulas of the different indexes used, we noted that Fulton's and Bjorndal's indexes were similar in their values; therefore, we considered these as one (Fulton's/Bjorndal index, hereafter). Additionally, simple linear regressions were used to analyze the relationship between the BC indexes and SCL, as well as between the BC indexes and W. Finally, a within-group sum of squares (WGSS) test was used to evaluate the BC classification of the specimens evaluated. Results were considered significant when $P < 0.05$. Mean and standard deviation (SD) are presented in the following notation: mean \pm SD. Analyses were performed using Past 4.08 statistical software (Hammer et al. 2001).

RESULTS


A total of 447 nesting olive ridley female turtles were sampled during the nesting season. The mean, minimum, and maximum values of the variables (SCL, SCW, TC, CCL, CCW, W) are presented in Table 1. The size (SCL, SCW) and W of nesting turtles was not significantly different from a normal distribution

(Shapiro Wilks test $SCL = 0.906$, $P = 0.0643$; Shapiro Wilks test $CCL = 0.858$, $P = 0.0003$; Shapiro Wilks test $w_{gt} = 0.995$, $P = 0.0003$; Fig. 2). A mean of 60.41 ± 2.98 cm was found for SCL (range = 32.7-71.1, $n = 450$), 52.69 ± 2.57 cm for SCW (range = 36-65.9, $n = 450$), and 34.35 ± 3.94 kg (range = 23.7-46.3, $n = 483$) for W. There was a positive relationship between W, and all measurements (Table 2), and particularly, a positive relationship was found between W and Cir (Fig. 3).

Thirty-three females were discarded from the BC analysis due to a failure in the digital scale during its sampling. The other 52 females were discarded from the BMI analysis due to the absence of thickness of carapace measures. The mean \pm SD Fulton's/Bjorndal body condition index estimated for the nesting olive ridley was 1.54 ± 0.15 , whereas the mean \pm SD of BMI was 2.18 ± 0.23 (Table 3). There was a significant difference between Fulton's/Bjorndal index and BMI (Mann-Whitney test = 0.0014, $P < 0.000001$). Moreover, there was a negative relationship between Fulton's/Bjorndal index and SCL ($r = -0.47$, $P < 0.000000783$; Fig. 4a), although there was a positive relationship between BMI and SCL ($r = 0.130$, $P < 0.004$; Fig. 4b). On the other hand, there was a positive

Table 1. Morphometric measurements of nesting female *L. olivacea* in La Escobilla, Oaxaca. SD: standard deviation, n: sample size.

Measurement	Abbreviation	Mean	SD	Range	n
Straight carapace length (cm)	SCL	60.41	2.98	32.7-71.1	450
Straight carapace width (cm)	SCW	52.69	2.57	36-65.9	450
Thickness of carapace (cm)	TC	25.65	2.07	13-43.9	399
Curved carapace length (cm)	CCL	65.11	3.33	57.4-93.7	483
Curved carapace width (cm)	CCW	69.17	3.43	60.4-98.2	483
Circumference (cm)	Cir	125.13	4.95	112.3-141.1	483
Weight (kg)	W	34.35	3.94	23.7-43.3	483

**Figure 2.** Size distribution for *L. olivacea* sea turtles in La Escobilla, Oaxaca. a) Straight carapace length, b) straight carapace width, c) weight. The line shows the normal distribution curve.

relationship between Fulton's/Bjorndal index and W ($r = 0.334$, $P < 0.000404$; Fig. 5a), and a positive relationship between BMI and W ($r = 0.517$, $P < 0.00005$; Fig. 5b).

Based on the Fulton's/Bjorndal index results, we classified 49 individuals with bad condition, 318 with good condition, and 47 with excellent condition. Using WGSS analysis, we found that 67.9% (WGSS = 41026, $F = 2.11$) of the individuals were correctly classified. Additionally, there were differences in the BCI index between BC groupings (Mann-Whitney test = 0.0014, $P < 0.000001$; Fig. 6a). On the other hand, BC charac-

terization based on BMI resulted in 28 individuals classified as in bad condition, 315 as in good condition, and 19 in excellent BC. There were differences in the BCI scores between the groups (Mann-Whitney test = 0.00001, $P < 0.0000001$; Fig. 6b), and using WGSS analysis, we found that 72.8% (WGSS = 2.14) of the individuals were correctly classified.

DISCUSSION

Morphometric studies provide basic information about animal development, evolution, biomechanics, behavior,

Table 2. Simple linear regression equations ($Y = ax + B$) for significant correlations of length measurements of *L. olivacea* in La Escobilla, Oaxaca. SCL: straight carapace length, SCW: straight carapace width, CCL: curved carapace length, CCW: curved carapace width, Cir: circumference, W: weight, r^2 : R-squared, F: F-test value, df: degrees of freedom, P : probability value (< 0.05).

X	Y	a	b	n	r^2	F	df	P
W	SCL	1.13	-34.25	443	0.54	22.34	1.29	0.0001
W	SCW	0.85	-11.24	443	0.28	12.85	1.05	0.0001
W	CCL	0.82	-19.58	443	0.44	18.94	1.11	0.0001
W	CCW	0.72	-16.18	443	0.36	15.95	0.97	0.0001
W	Cir	0.72	-16.18	443	0.36	15.95	0.97	0.0001




Figure 3. Relationship between circumference and weight of nesting female *L. olivacea* in La Escobilla, Oaxaca.

Table 3. Body condition index comparisons of three formulae applied in *L. olivacea* in La Escobilla, Oaxaca. SD: standard deviation, n: sample size.

	Mean	SD	Range	n
Fulton index	1.54	0.15	0.88-2.59	414
Bjorndal index	1.54	0.15	0.88-2.59	414
BM Index	2.18	0.20	1.18-4.01	362

ecology, and physiology (van Dam & Diez 1998); furthermore, they play a key role in characterizing and analyzing intra-population trends (Figueroa & Alvarado 1990, van Dam & Diez 1998). For the olive ridley turtle, knowledge about their morphometric characteristics and BC has remained limited. In this study, BC was classified using morphometric analyses, and it is the first assessment for olive ridleys nesting in La Escobilla on the Pacific coast of southern Mexico. It also shows that fewer nesting females were registered at this location a few decades later.

Body size is commonly used in field studies with sea turtles; nonetheless, one of the most common body measures is the curve measure (Sönmez 2019). Here, we report two groups of measures (straight and curved). Our mean SCL (60.41 cm) and SCW (52.69 cm) found were smaller than SCL (62.98 cm) and SCW (57.07 cm) reported by Frazier (1983) ($n = 82$), also on nesting olive ridley turtles in La Escobilla. More recently, Gaona & Barragán (2016) reported a range of CCL from 60 to 78 cm, a range of CCW from 63 to 78 cm, and a range of W from 33 to 52 kg. Conversely, the mean results found in this study were smaller than those of the previous studies mentioned above. Espinoza-Romo et al. (2018) reported a mean SCL of 61.7 ± 1.5 and a mean W of 34.9 ± 3.7 for adult olive ridley turtles in northern Sinaloa, Mexico. However, in this study, they evaluated turtles captured while they floated at the surface of the open sea, including data from adult males.




Figure 4. Relationship between a) straight carapace length (cm) and Fulton's/Bjorndal index, and b) straight carapace length (cm) and body mass index (BMI) in La Escobilla, Oaxaca.




Figure 5. Relationship between a) weight and Fulton's/Bjorndal index, and b) weight and body mass index BMI in La Escobilla, Oaxaca.

One of the advantages of getting body measurements from turtles is that regional differences or latitudinal trends can be evaluated across nesting populations (Tiwari & Bjorndal 2000). For example, the CCL and CCW (65.11 ± 3.33 ; 69.17 ± 3.43 cm, respectively) obtained were smaller than those reported by Kalb (1999) (CCL: $68.4 \text{ cm} \pm 0.2 \text{ cm}$) for Playa Nancite, Costa Rica, and Vera & Rosales (2012) in Tumbes, Peru (CCL: $67.8 \text{ cm} \pm 2.9 \text{ cm}$). Dornfeld et al. (2015) reported a mean CCL of $65.9 \pm 3.5 \text{ cm}$ for Playa Grande, Costa Rica, while in the western Pacific, Whiting et al. (2007) reported a mean CCL of $69.6 \pm 2.3 \text{ cm}$. In the western Atlantic, da Silva et al. (2007) reported a mean CCL of $73.1 \pm 0.2 \text{ cm}$ for Sergipe and Bahía, Brazil. Although our results are slightly smaller than those in the literature, Bergmann's rule proposes a positive relationship between mean body size and latitude, in which smaller individuals are found at lower

latitudes (Gardner et al. 2011, Angielczyk et al. 2015); however, this is only applied for endothermic animals (Ashton & Feldman 2003).

Some hypotheses proposed by Le Gouvello et al. (2020a) have been used to explain the mechanisms responsible for a decline in sea turtle mean carapace size. First, the possibility of an increase in the number of first-time nesters in the nesting population, which means an increase in food competition and consequently affects growth rates (density-dependence processes). Second, a change in the size at maturity, with females reaching maturity at a smaller size. Third, lower rate of post-maturity growth, and fourth, decreased adult survivorship. However, a fifth hypothesis related to ecosystem variation productivity in tropical regions has been proposed as a product of large-scale ocean-atmosphere anomalies affecting primary and secondary production. In the west Atlantic,




Figure 6. Body condition scores of nesting females of *L. olivacea* in La Escobilla, Oaxaca. a) Fulton's/Bjorndal index, b) body mass index (BMI).

Bjorndal et al. (2017) showed that ecological regime shifts due to the synergistic effect of a strong El Niño-Southern Oscillation and the intensification of the warming rate over the last two to three decades resulted in decreased growth rates of green turtles. As a consequence of any of these hypotheses, smaller females may produce smaller hatchlings of lower fitness (Le Gouvello et al. 2020b). In addition, smaller females may produce fewer eggs, affecting overall population growth (Le Gouvello et al. 2020a).

Regarding the relationships between body measurements and W , positive correlations were found in most cases, the strongest being between Cir and W . In this sense, Cir explains 61% of the increment in W or vice-versa. Although this is an interesting result, there are no similar studies, perhaps because mass measurements may be difficult to obtain, particularly for large animals in challenging field situations (Thomson et al. 2009). Weighing large sea turtles along nesting beaches in remote field sites is logistically difficult, and it often involves transportation of the animals to weighing facilities and holding them for prolonged periods (Thomson et al. 2009), which may affect mass measurements.

BC is a proxy for a turtle's health condition. It can be quantified by assessing an animal's external parameters (morphometrics measures) (Bjorndal et al. 2000, Labrada-Martagon et al. 2010), which is a reference to its energetic condition since an individual in good condition is assumed to have higher energy reserves than one in poor condition (Labrada-Martagon et al. 2010). Page-Karjian & Perrault (2021) explained that the first step to assessing the health of sea turtle populations is to generate population-specific baseline health data to establish a health database representative for that population. Our study assessed the BC for olive

ridley nesting at La Escobilla. BC is correlated with fecundity because energetic reserves limit the amount of energy that can be distributed for reproduction and can supply valuable information about overall fitness and health (Perrault et al. 2012). This information can be especially relevant for threatened populations to prioritize conservation actions (Schulte-Hostedde et al. 2005). Sea turtles can be sensitive to the decrease in their BC because of their migratory behavior and the fact that they do not reproduce annually (Bjorndal et al. 2003). Moreover, females concentrate their reproductive efforts in specific years, followed by non-nesting years, reducing the number of breeding migrations between their distant nesting and feeding grounds (Bjorndal et al. 2000, Marco et al. 2011).

Health indicators, including BC and physical examination, have been used to determine the general health status of multiple free-living species (Thomson et al. 2009). Our study found similar BC values - using Fulton's/Bjorndal index as a reference - to those reported in healthy populations of olive ridley turtles (*L. olivacea*) in northern Sinaloa, Mexico (Espinoza-Romo et al. 2018), and was greater than those reported in Brazil (de Deus et al. 2015) and Venezuela (Barrios et al. 2015) where most turtles had a poor BC and included individuals with signs of physical illness and disease.

This study compared two ways to estimate the BC and another index (BMI) commonly implemented in veterinary sciences and cattle ranching. The BMI presented here allowed us to include the accumulation of a large fat area in the turtles' internal cavities. Therefore, BMI can be an alternative tool to predict BC in sea turtles because fat plays an important role in several functions (e.g. source of energy, buoyancy, thermal insulator) (Young 1976).

Although visual assessment techniques of the physical condition have been proposed in field studies for wildlife health assessments (Thomson et al. 2009), they are inadequate in identifying BC scores; hence, using the BC index is adequate. The BC index and the categories established (i.e. bad, good, and excellent condition) represent a viable alternative to assign each sea turtle with a BC class, and based on the percentages of the WGSS test, this classification is straightforward. Thus, similar to Stevenson & Woods (2006), we suggest that using different indexes to evaluate BC in the same population will reflect the status of the population.

In a changing ocean environment, it is important to provide baseline observations on ecosystems and organisms' health (Nolte et al. 2020). This study is the first to report the smallest female size of nesting olive ridleys for the southwest Mexican Pacific coast and is the first estimation of BC showing that most of the nesting females in La Escobilla have good body condition; however further research is required to understand if smaller female sizes can affect the reproductive output and overall population growth.

ACKNOWLEDGMENTS

We thank the Universidad Autónoma del Estado de México (UAEMex) and its Division of Postgraduate Studies (PCARN) for the logistics and facilities provided. To the Universidad del Mar for the facilities provided through the project CUP: 2IR2104. This work was carried out under permits SGPA/DGVS/03919/21 from Secretaria del Medio Ambiente y Recursos Naturales (for handling and extracting blood). Special thanks go to Iain Sinclair for his revisions and suggestions that improved the English of this manuscript. ABS thanks CONACyT for the scholarship grant. JG-G, PS-N, and MLRG thank the Sistema Nacional de Investigadores (SNI) for their recognition and support. This paper is part of the Ph.D. dissertation of ABS as a student of the Division of Postgraduate Studies (PCARN) at UAEMex.

REFERENCES

- Abreu-Grobois, A. & Plotkin, P. 2008. *Lepidochelys olivacea*. In: The IUCN Red List of Threatened Species. IUCN. doi: 10.2305/IUCN.UK.2008.RLTS.T11534A3292503
- Angielczyk, K.D., Burroughs, R.W. & Feldman, C.R. 2015. Do turtles follow the rules? Latitudinal gradients in species richness, body size, and geographic range area of the world's turtles. *Journal of Experimental Zoology*, 324: 270-294. doi: 10.1002/jez.b.22602
- Ashton, K.G. & Feldman, C.R. 2003. Bergmann's rule in non-avian reptiles: turtles follow it, lizards and snakes reverse it. *Evolution*, 57: 1151-1163. doi: 10.1111/j.0014-3820.2003.tb00324.x
- Barrios, H., Espinoza, N., Shima, T. & Wildermann, N.E. 2015. Body condition index in rescued green turtles (*Chelonia mydas*) in the Gulf of Venezuela: a seven-year assessment. In: 35th Annual Symposium on Sea Turtle Biology and Conservation, MACART Press, Turkey.
- Barton, B.A., Morgan, J.D. & Vijayan, M.M. 2002. Physiological and condition-related indicators of environmental stress in fish. In: Adams, S.M. (Ed.). *Biological indicators of aquatic ecosystem stress*. American Fisheries Society, Bethesda, pp. 111-148.
- Bernardon, J. & Plotkin, P.T. 2007. An evolutionary perspective on the arribada phenomenon and reproductive behavioural polymorphism of olive ridley sea turtles (*Lepidochelys olivacea*). In: Plotkin, P.T. (Ed.). *Biology and conservation of ridley sea turtles*. John Hopkins University Press, Maryland, pp. 59-87.
- Beverton, R.J.H. & Holt, S.J. 1957. *On the dynamics of exploited fish population*. Chapman & Hall Press, London.
- Bézy, V.S., Girondot, M. & Valverde, R.A. 2016. Estimation of the net nesting effort of olive ridley arribada sea turtles based on nest densities at Ostional beach, Costa Rica. *Journal of Herpetology*, 50: 409-415. doi: 10.1670/14-152
- Bjorndal, K., Bolten, A.B. & Chaloupka, M.Y. 2000. Green turtle somatic growth model: evidence for density dependence. *Ecological Applications*, 10: 269-282.
- Bjorndal, K.A., Bolten, A.B. & Chaloupka, M.Y. 2003. Survival probability estimates for immature green turtles, *Chelonia mydas*, in the Bahamas. *Marine Ecology Progress Series*, 252: 273-281.
- Bjorndal, K.A., Bolten, A.B., Chaloupka, M., Saba, V., Bellini, C., Marcovaldi, M.A.G., et al. 2017. Ecological regime shift drives declining growth rates of sea turtles throughout the West Atlantic. *Global Change Biology*, 23: 4556-4568. doi: 10.1111/gcb.13712
- Campbell, L.M. 1998. Use them or lose them? Conservation and the consumptive use of marine turtle eggs at Ostional, Costa Rica. *Environment Conservation*, 25: 305-319. doi: 10.1017/S0376892998000393

- Campbell, L.M. 2007. Understanding human use of olive ridleys, implications for conservation. In: Plotkin, P.T. (Ed.). *Biology and conservation of ridley sea turtles*. Johns Hopkins University Press, Maryland, pp. 23-43.
- Castro, M.I., Auriolos, D., Montaña, S., Pérez, F. & López, N. 2001. Lípidos totales, colesterol y triglicéridos en crías de lobo marino de California del Golfo de California. *Ciencias Marinas*, 27: 375-396.
- Cone, R.S. 1989. The need to reconsider the use of conditions indices in fishery science. *Transactions of the American Fisheries Society*, 118: 510-514.
- Copeland, T. 2004. An evaluation of relative weight as an indicator of body composition and nutritional status in wild fish. Doctoral Dissertation, Virginia Tech, Virginia.
- Cornelius, S.E. 1986. The sea turtles of Santa Rosa National Park, Costa Rica. *Fundación de Parques Nacionales*, San José.
- Cornelius, S.E., Ulloa, M.A., Castro, J.C., Mata, M. & Robinson, D.C. 1991. Management of olive ridley sea turtles (*Lepidochelys olivacea*) nesting at Playas Nancite and Ostional, Costa Rica. In: Robinson, J.G. & Redford, K.H. (Eds.). *Neotropical wildlife use and conservation*. The University of Chicago Press, Chicago, pp. 111-135.
- Cornelius, S.E., Arauz, R., Fretey, J., Godfrey, M.H., Márquez, R. & Shanker, K. 2007. Effect of land-based harvest of *Lepidochelys olivacea*. In: Plotkin, P.T. (Ed.). *Biology and conservation of ridley sea turtles*. Johns Hopkins University Press, Maryland, pp. 231-251.
- da Silva, A.C., de Castilhos, J.C., Lopez, G.G. & Barata P.C. 2007. Nesting biology and conservation of the olive ridley sea turtle (*Lepidochelys olivacea*) in Brazil, 1991/1992 to 2002/2003. *Journal of Marine Biology Association of the United Kingdom*, 87: 1047-1056. doi: 10.1017/S0025315407056378
- de Deus, M.R., Silva, A., Baptistotte, C. & Work, T.M. 2015. Health condition of juvenile *Chelonia mydas* related to fibropapillomatosis in southeast Brazil. *Diseases of Aquatic Organisms*, 115: 193-201. doi: 10.3354/dao02883
- Deem, S.L., Norton, T.M., Mitchell, M., Segars, A., Alleman, A.R. & Cray, C. 2009. Comparisons of blood values in foraging, nesting, and stranded loggerhead turtles (*Caretta caretta*) along the coast of Georgia, USA. *Journal of Wildlife Diseases*, 45: 41-56. doi: 10.7589/0090-3558-45.1.41
- Diario Oficial de la Federación (DOF). 2010. Norma Oficial Mexicana NOM-059-SEMARNAT-2010, que determinan las especies de flora y fauna silvestres, terrestres y acuáticas, endémicas, amenazadas, en peligro de extinción y sujetas a protección especial. Gobierno Federal, Ciudad de México.
- Dornfeld, T.C., Robinson, N.J., Santidrián, P. & Paladino, F. 2015. Ecology of solitary nesting olive ridley sea turtle at Playa Grande, Costa Rica. *Marine Biology*, 162: 123-139. doi: 10.1007/s00227-014-2583-7
- Espinoza-Romo, B.A., Sainz-Hernández, J.C., Ley-Quinónez, C.P., Hart, C.E., Leal-Moreno, R., Aguirre, A.A., et al. 2018. Blood biochemistry of olive ridley (*Lepidochelys olivacea*) sea turtles foraging in northern Sinaloa, Mexico. *Plos One*, 13: e0199825. doi: 10.1371/journal.pone.0199825
- Figuroa, A. & Alvarado, J. 1990. Morphometric comparison of the *Chelonia* populations of Michoacan, Mexico, and Tortuguero, Costa Rica. In: Epperly, S.P. & Braun, J. (Eds.). *Proceedings of the tenth annual workshop on sea turtle biology and conservation*. NOAA Technical Memorandum NMFS-SEFC-278, USA.
- Flint, M., Morton, J.M., Limpus, C.J., Patterson-Kane, J.C., Murray, P.J. & Mills, P.C. 2010. Development and application of biochemical and hematological reference intervals to identify unhealthy green sea turtles (*Chelonia mydas*). *Veterinary Journal*, 185: 299-304. doi: 10.1016/j.tvjl.2009.06.011
- Fonseca, L.G., Murillo, G.A., Guadamúz, L., Spínola, R.M. & Valderde, R.A. 2009. Downward but stable trend in the abundance of arribada olive ridley sea turtles (*Lepidochelys olivacea*) at Nancite beach, Costa Rica (1971-2007). *Chelonian Conservation Biology*, 8: 19-27.
- Frazier, J. 1983. Misidentification of sea turtles in the east Pacific: *Caretta* and *Lepidochelys olivacea*. *Journal of Herpetology*, 19: 1-11. doi: 10.2307/15644114
- Gaona, O. & Barragán, A.R. 2016. Las tortugas marinas en México: logros y perspectivas para su conservación. *Comisión Nacional de Áreas Naturales Protegidas*, Ciudad México.
- García, E. 1981. Modificaciones al sistema de clasificación climática de Köppen para adaptarlo a las condiciones de la República Mexicana. *Universidad Nacional Autónoma de México*, Ciudad de México.
- Gardner, J.L., Peters, A., Kearny, M.R., Joseph, L. & Heinsohn, R. 2011. Declining body size: a third universal response to warming? *Trends in Ecology & Evolution*, 26: 285-291. doi: 10.1016/j.tree.2011.03.005
- Green, A.J. 2001. Mass/length residuals: Measures of body condition or generators of spurious results?

- Ecology, 82: 1473-1483. doi: 10.1890/00129658(2001)082[1473:MLRMOB]2.0.CO;2
- Hammer, Ø., Harper, D.A.T. & Ryan, P.D. 2001. PAST: Paleontological statistics software package for education and data analysis. *Palaeontologia Electronica*, 4: 1-9.
- Hays, G.C. & Speakman, J.R. 1991. Reproductive investment and optimum clutch size of loggerhead sea turtles (*Caretta caretta*). *Journal of Animal Ecology*, 60: 455-462.
- Hope, R.A. 2002. Wildlife harvesting, conservation and poverty: The economics of olive ridley egg exploitation. *Environmental Conservation*, 29: 375-384.
- Hornarvar, S., O'Connor, M.P. & Spotila, J.R. 2008. Density-dependent effects on hatching success of the olive ridley turtle, *Lepidochelys olivacea*. *Oecologia*, 157: 221-230. doi: 10.1007/s00442-008-1065-3
- Ilgaz, Ç., Türkozan, O., Özdemir, A., Kaska, Y. & Stachowitsch, M. 2007. Population decline of loggerhead turtles: two potential scenarios for Fethiye beach, Turkey. *Biological Conservation*, 16: 1027-1037.
- Jessop, T.S., Summer, J.M., Limpus, C.J. & Whitakker, J.M. 2004. Interplay between plasma hormone profiles, sex, and body condition in immature hawksbill turtles (*Eretmochelys imbricata*) subjected to a capture stress protocol. *Comparative Biochemistry and Physiology - Part A: Molecular & Integrative Physiology*, 137: 197-204. doi: 10.1016/j.cbpa.2003.09.029
- Jones, R.E., Petrell, R.J. & Pauly, D. 1999. Using modified length-weight relationships to assess the condition of fish. *Aquaculture Engineering*, 20: 261-276. doi: 10.1016/S0144-8609(99)00020-
- Kalb, H.J. 1999. During the interesting period, the behavior and physiology of solitary and arribada nesting olive ridley sea turtles (*Lepidochelys olivacea*). Ph.D. Thesis, Wittenberg University, Ohio.
- Kophamel, S., Ward, L.C., Mendez, D., Ariel, E., Bell, I., Shum, E., et al. 2023. Adipose tissue estimation of foraging and nesting green turtles *Chelonia mydas* using bioelectrical impedance analysis. *Endangered Species Research*, 51: 127-142. doi: 10.3354/esr01248
- Labrada-Martagon, V., Méndez-Rodríguez, L.C., Gardner, S., Cruz-Escalona, V. & Zenteno, T. 2010. Health indices of green turtle (*Chelonia mydas*) along the Pacific of Baja California Sur, Mexico. II. Body condition index. *Chelonian Conservation and Biology*, 9: 173-183. doi: 10.2744/CCB-0807.1
- Lamont, M.M. & Johnson, D. 2021. Variation in species composition size and fitness of two multi-species sea turtle assemblages using different neritic habitats. *Frontiers in Marine Science*, 7: 608740. doi: 10.3389/fmars.2020.608740
- LeBlanc, A., Rostal, D.C., Drake, K.K., Williams, K.L., Robinette, J. & Barnard-Keinath, D.E. 2014. The influence of maternal size on the eggs and hatchlings of loggerhead sea turtles. *Southeast Naturalist*, 13: 587-599.
- LeCren, E.D. 1951. The length-weight relationship and seasonal cycle in gonad weight and condition in the perch (*Perca fluviatilis*). *Journal of Animal Ecology*, 20: 201-219.
- Le Gouvello, D., Nel, R. & Cloete, A. 2020b. The influence of individual size on clutch size and hatchling fitness traits in sea turtles. *Journal of Experimental Marine Biology and Ecology*, 527: 151372.
- Le Gouvello, D., Girondot, M., Bachoo, S. & Nel, R. 2020a. The good and bad news of long-term monitoring: an increase in abundance but decreased body size suggests reduced potential fitness in nesting sea turtles. *Marine Biology*, 167: 112.
- Marco, A., Abella-Pérez, E., Monzón, C., Martins, S., Araújo, C. & López, L.F. 2011. The international importance of the Archipelago of Cape Verde for marine turtles, in particular the loggerhead turtle *Caretta*. *Zoologia Caboverdiana*, 2: 1-11.
- Márquez, R., Peñaflores, C. & Vasconcelos, J. 1996. Olive ridley turtles (*Lepidochelys olivacea*) show signs of recovery at La Escobilla, Oaxaca. *Marine Turtle Newsletter*, 73: 5-7.
- Márquez, R., Jiménez-Quiroz, M.C., Peñaflores, C. & Garduño, M. 2014. México y las tortugas marinas. In: Márquez, R. & Garduño, M. (Eds.). *Tortugas marinas*. Instituto Nacional de la Pesca, Ciudad de México, pp. 13-37.
- Mazaris, A.D., Schofield, G., Ghazinou, C., Almpnidou, V. & Hays, G.C. 2017. Global sea turtle conservation successes. *Marine Conservation*, 3: e1600730.
- Meffe, G.K. 1999. Conservation medicine. *Conservation Biology*, 13: 953-954.
- Munson, L. & Karesh, W.B. 2002. Disease monitoring for the conservation of terrestrial animals. In: Aguirre, A.A., Ostfeld, R.S., Tabor, G.M., House, C. & Pearl, M.C. (Eds.). *Conservation medicine, ecological health in practice*. Oxford University Press, New York, pp. 95-103.
- Nájera-Hillman, E., Bass, J.B. & Buckham, S. 2012. Distribution patterns of the barnacle, *Chelonibia testudinaria*, on juvenile green turtles (*Chelonia mydas*) in Bahía Magdalena, Mexico. *Revista*

- Mexicana de Biodiversidad, 83: 1171-1179. doi: 10.7550/rmb.27444
- Nel, R., Punt, A.E. & Hughes, G.R. 2013. Are coastal protected areas always effective in achieving population recovery for nesting sea turtles? Plos One, 8: e63525. doi: 10.1371/journal.pone.006.3525
- Nolte, C.R., Nel, R. & Pfaff, M.C. 2020. Determining body condition of nesting loggerhead sea turtles (*Caretta caretta*) in the southwest Indian Ocean. Journal of the Marine Biological Association of the United Kingdom, 100: 1-9. doi: 10.1017/S0025315420000107
- Ocana, M., Harfush, M. & Hepell, S.S. 2012. Mass nesting of olive ridley sea turtles *Lepidochelys olivacea* at La Escobilla, Mexico: Linking nest density and rates of destruction. Endangered Species Research, 16: 45-54. doi: 10.3354/esr00388
- Page-Karjian, A. & Perrault, J. 2021. Sea turtle health assessments: Maximizing turtle encounters to better understand health. In: Nahill, B. (Ed.). Sea turtle research and conservation. Lessons from working in the field. Academic Press, London, pp. 31-44.
- Peig, J. & Green, A.J. 2010. The paradigm of body condition: a critical reappraisal of current methods based on mass and length. Functional Ecology, 24: 1323-1332. doi: 10.1111/j.1365-2435.2010.01751.x
- Peñaflores, C.J., Vasconcelos, J. & Albavera-Márquez, E. 2000. Twenty-five years nesting of olive ridley sea turtle *Lepidochelys olivacea* in Escobilla beach, Oaxaca, México. In: Grobois, F.A., Briseño-Dueñas, R. & Sarti, L. (Eds.). Proceedings of 18th International Sea Turtle Symposium. NOAA Technical Memorandum NMFS-SEFSC-436, USA, pp. 27-29.
- Pérez-Castañeda, R., Salum-Fares, A. & Defeo, O. 2007. Reproductive patterns of the hawksbill turtle *Eretmochelys imbricata* in sandy beaches of the Yucatan Peninsula. Journal of the Marine Biological Association of the United Kingdom, 87: 815-824.
- Perrault, J., Miller, D.L., Eads, E., Johnson, C., Merrill, A., Thompson, L.J., et al. 2012. Maternal health status correlates with nest success of leatherback sea turtles (*Dermochelys coriacea*) from Florida. Plos One, 7: e31841. doi: 10.1371/journal.pone.0031841
- Phasey, H., Roberts, D.L., Rojas-Cañizales, D., Mejías-Balsalobre, C., Griffiths, R.A. & Williams-Guillen, K. 2020. Using GPS-enabled decoy turtle eggs to track illegal trade. Current Biology, 30: 1066-1068. doi: 10.1016/j.cub.2020.08.065
- Piacenza, S.E., Balazs, G.H., Hargrove, S.K., Richards, P.M. & Hepell, S.S. 2016. Trends and variability in demographic indicators of a recovering population of green sea turtles *Chelonia mydas*. Endangered Species Research, 31: 103-117.
- Plotkin, P.T., Rostal, D.C., Byles, R.A. & Owens, D.W. 1997. Reproductive and developmental synchrony in female *Lepidochelys olivacea*. Journal of Herpetology, 31: 17-22. doi: 10.2307/1565233
- Ramírez-Villanueva, R.I., Gumeta-Gómez, F., Lara-Uc, M., López-Vivas, J.M. & Hinojosa-Arango, G. 2023. The use of the territorial sea by *Lepidochelys olivacea* (Eschscholtz, 1829) in front of nesting beaches in Oaxaca, Mexico. Regional Studies in Marine Science, 65: 103065.
- Resendiz, E., Fernández, H. & Lara-Uc, M.M. 2018. Baseline health indicators of eastern Pacific green turtles (*Chelonia mydas*) from Baja California Sur, Mexico. Comparative Clinical Pathology, 27: 1309-1320. doi: 10.1007/s00580-0182740-3
- Rossi, S., Sánchez, A.M., Guimaraes, R., Ramblas, R., Setim, F., Gattamnorta, M., et al. 2019. Monitoring green sea turtles in Brazilian feeding areas: relating body condition index to fibropapillomatosis prevalence. Journal of Marine Biological Association of the United Kingdom, 99: 1879-1887. doi: 10.1017/S0025315419000730
- Salazar-Cuytún, E., Sarmiento, L.A., Aguilar, A.J., Fonseca, M. & Tedeschi, L. 2020. Body mass index and body chemical components in Pelibuey ewes. Ecosistemas y Recursos Tropicales, 7: e2515. doi: 10.19136.era.a7n2.2515
- Schulte-Hostedde, A.L., Zinner, B., Millar, J.S. & Hickling, G.J. 2005. Restitution of mass-size residuals: Validating body condition indices. Ecology, 86: 155-163. doi: 10.1890/04-0232
- Seminoff, J.A., Reséndiz-Hidalgo, A., Jiménez de Reséndiz, B., Wallace, J.N. & Todd-Jones, T. 2008. Tortuga marinas. In: Danemann, G.D. & Ezcurra, E. (Eds.). Bahía de los Ángeles: recursos naturales y comunidad. Secretaría de Medio Ambiente y Recursos Naturales, Ciudad de México, pp. 457-493.
- Shanker, K., Pandav, B. & Choudhury, B.C. 2003. An assessment of the olive ridley turtle (*Lepidochelys olivacea*) nesting population in Orissa, India. Biological Conservation, 115: 149-160.
- Sönmez, B. 2019. Morphological variations in the green turtle (*Chelonia mydas*): a field study on an Eastern Mediterranean nesting population. Zoological Studies, 58: 1-13. doi: 10.6620/ZS.2019.58-16
- Stamper, M.A., Harms, C., Epperly, S.P., Braun, J., Avens, L. & Stoskopf, M.K. 2005. Relationships between barnacle epibiotic load and hematologic parameters in loggerhead sea turtles (*Caretta caretta*),

- a comparison between migratory and residential animals in Pamlico Sound, North Carolina. *Journal of Zoo and Wildlife Medicine*, 36: 635-641. doi: 10.1638/04-074.1
- Stevenson, R. & Woods, W.A. 2006. Condition indices for conservation: new uses for evolving tools. *Integrative and Comparative Biology*, 46: 1169-1190. doi: 10.1093/icb/icl052
- Steward, A.Y. 2001. Poached modernity: Parks, people and politics in Nicaragua, 1975-2000. Ph.D. Dissertation, Rutgers University, New Jersey.
- Taylor, J.A. 1979. The foods and feeding habits of subadult *Crocodylus porosus* Schneider in northern Australia. *Australian Wildlife Research*, 6: 347-359.
- Thomson, J., Burkholder, D., Haithaus, M.R. & Dill, L. 2009. Validation of a rapid visual-assessment technique for categorizing the body condition of green turtles (*Chelonia mydas*) in the field. *Copeia*, 2: 251-255. doi: 10.1643/CE-07-227
- Tiwari, M. & Bjorndal, A.K. 2000. Variation in morphology and reproduction in loggerheads, *Caretta caretta*, nesting in the United States, Brazil, and Greece. *Herpetologica*, 56: 343-356.
- Tuček, J.B., Nel, R., Girondot, M. & Hughes, G. 2014. Age-size relationship at reproduction of South African female loggerhead turtles *Caretta caretta*. *Endangered Species Research*, 23: 167-175.
- Ullman-Culleré, M.H. & Foltz, C.J. 1999. Body condition scoring: a rapid and accurate method for assessing health status in mice. *Comparative Medicine*, 49: 319-323.
- Valverde, R.A., Orrego, C.M., Tordoir, M.T., Gómez, F., Solís, D., Hernández, R., et al. 2012. Olive ridley mass nesting ecology and egg harvest at Ostional beach, Costa Rica. *Chelonian Conservation and Biology*, 11: 1-11.
- van Dam, R.P. & Diez, C.E. 1998. Caribbean hawksbill turtles morphometrics. *Bulletin of Marine Sciences*, 62: 145-155.
- Vera, M. & Rosales, C.A. 2012. Estructura de tallas de tortugas pico de loro *Lepidochelys olivacea* (Testudines: Cheloniidae) en Tumbes, Perú. *Revista Peruana de Biología*, 19: 175-180.
- Whiting, S., Long, J.L., Hadden, K.M., Anderson, D.K. & Koch, A. 2007. Insights into size, seasonality, and biology of a nesting population of the olive ridley turtle in northern Australia. *Wildlife Research*, 34: 200-210. doi: 10.1071/WR06131
- Young, R.A. 1976. Fat, energy, and mammalian survival. *American Zoologist*, 16: 699-710. doi: 10.1093/icb/16.4.699

Received: August 17, 2023; Accepted: June 13, 2024